


Hallux Metatarsophalangeal Joint Fusion



This operation is intended to glue (fuse) the big toe joint together and is usually performed as a day case operation. It is also called an arthrodesis of the big toe joint (metatarsophalangeal joint).

It is usually performed for painful arthritis of this joint (hallux rigidus) which has not responded to other non-operative treatment. The joint surfaces are removed via an incision over the upper surface or inside surface of the foot. The two 'raw' bone surfaces are then applied together and screws used to compress them together and stabilise the joint in the desired position. These screws do not usually need to be removed subsequently although they can be if need be. Once fused, the big toe does not move except at the end joint (interphalangeal joint) and this should greatly improve / abolish the painful symptoms from this joint. The toe is positioned according to the need to roll off the big toe, exercise, or wear higher heeled shoes. Once fusion is successful, you will be able to wear most shoes (but not necessarily all types). There are no limits to exercise activities after the fusion.

General Recovery Facts

- Operation performed under general anaesthetic or regional anaesthetic
- You are able to walk on the heel of the foot the day of surgery
- You must wear your surgical shoe (heel wedge shoe) at all times including in bed at night
- You may not walk on the foot at all even in the house without this shoe
- You may not drive after the surgery for six weeks unless you have an automatic vehicle and only the left foot has undergone surgery
- The surgical shoe is worn for 6-8 weeks

Main Risks of Surgery

Swelling - Initially the foot will be very swollen and needs elevating. The swelling will disperse over the following weeks and months but will be apparent for up to 6-9 months.

Infection - This is the biggest risk with this type of surgery. Smoking increases the risk 16 times. You will be given intravenous antibiotics to help prevention. However, the best way to reduce your chances of acquiring an infection is to keep the foot elevated over the first 10 days. If there is an infection, it may resolve with a course of antibiotics.

Wound problems - Sometimes the wounds can be slower to heal and this does not usually cause a problem but needs to be closely observed for any infection occurring.

Scar sensitivity - The scars can be quite sensitive following surgery but this usually subsides without treatment. If persistent sensitivity occurs then this can be treated.

Nerve Injury - The risk of the small nerves in the area being directly injured by the surgeon is approximately 1%. However, the nerves can become bruised by the surgery as a result of the swelling (10%). Whilst this usually recovers, you could end up with some permanent numbness over the big toe area, which might cause irritation.

CRPS - This stands for complex regional pain syndrome. It occurs rarely (1%) in a severe form and is not properly understood. It is thought to be inflammation of the nerves in the foot and it can also follow an injury. We do not know why it occurs. It causes swelling, sensitivity of the skin, stiffness and pain. It is treatable but in its more severe form can take many months to recover.

Delayed and non union - This is when the bones fail to join and bone has not grown across the cut bone. If this is painful then further surgery may be needed. The risk of this is approximately 10%.

Hallux Metatarsophalangeal Joint Fusion

Main Risks Of Surgery

Malposition - ideally the toe is fused in a position that allows optimum function and gives the best appearance. We will try to fuse the joint in the best position, using a flat panel to simulate the normal standing posture of the foot. However, as you are asleep and lying down, it is not always possible to achieve this best position. The toe may be either too high or too low. This rarely causes a problem that requires further surgery and can usually be accommodated with insoles.

Deep Vein Thrombosis (DVT) - This is a clot in the deep veins of the leg and the risk of this occurring following foot and ankle surgery is low (generally < 1%). The fact that you are mobile after surgery and able to take weight through the heel of the operated foot helps to minimise this small risk. However, it is sensible to try and move the toes and the ankle regularly following the surgery and probably also sensible to avoid a long-haul flight in the first 4 weeks following surgery. If a deep vein thrombosis (DVT) occurs then you will require treatment with heparin and Warfarin to try and prevent any of the clot travelling to the lungs (pulmonary embolus / PE) which can be much more serious).

Stiffness - The big toe joint is almost always more stiff following this surgery because of the scar tissue that forms. The stiffness can be minimised by beginning to move the big toe after 2 weeks from surgery and your surgeon will advise you regarding this. Most of the movement usually returns but some stiffness may remain permanently.

Continuing symptoms - Most people (90%) are very happy with the results of their bunion surgery but you can appreciate that if some of the above problems occur then this may affect the end result. Occasionally (5%) the bunion may recur although not usually to the same degree. This is more likely in patients who have more lax ligaments in their feet. If there is a recurrence then you don't necessarily require any further surgery - this will depend upon your symptoms.

Sick Leave

In general 2 weeks off work is required for sedentary employment, 6 weeks for work involving standing or walking, and 8 to 10 weeks for manual labour work. We will provide a sick certificate for the first 2 weeks; further notes can be obtained from your GP.

Driving

IF have an AUTOMATIC VEHICLE and ONLY LEFT leg surgery then it is likely you will be allowed to drive after your outpatient review at 2 weeks post surgery. IF you have a MANUAL VEHICLE or RIGHT leg surgery then you will NOT be able to drive until 6 weeks post surgery.

These notes are intended as a guide and some of the details may vary according to your individual surgery or because of special instructions from your surgeon.

sussexfoot&anklecentre

The Sussex Foot & Ankle Centre was founded in 2005 by two orthopaedic surgeons, David Redfern and Stephen Bendall, with the aim of providing a high quality specialist service for the diagnosis and treatment of all foot and ankle problems. Both orthopaedic surgeons are specialists in problems affecting the foot and ankle and have many years of experience. They operate the service with outpatient clinics at the Brighton and Haywards Heath Nuffield Hospitals.

The sussex foot and ankle center strives to provide the best advice and treatment for all foot and ankle problems. This includes sports injuries and trauma, bunions, metatarsalgia, and arthritis. Both surgeons have particular interests in minimally invasive surgery and are at the forefront of developing such techniques in this country.

Both surgeons are also academically very active and have appointments within the national (BOFAS) and international (EFAS) professional foot and ankle surgery societies.

David J Redfern
MBBS, FRCS, FRCS (Tr & Orth)
Consultant Orthopaedic Surgeon

Stephen P Bendall
MBBS, FRCS, FRCS (Orth)
Consultant Orthopaedic Surgeon

Private Secretary 01273 627060
Direct Fax 01273 627061
Email admin@sussexfoot.com

Haywards Heath
Sussex Nuffield Hospital
Burrell Road, Haywards Heath
East Sussex RH16 1UD
Tel 01444 456999

Brighton
Sussex Nuffield Hospital
Warren Road, Woodingdean
East Sussex BN2 6DX
Tel 01273 627060